

A new chiral electro-optic effect: Sum-frequency generation from optically active liquids in the presence of a dc electric field

P. Fischer,^{1,2,*} K. Beckwitt,² A.D. Buckingham,³ D.S. Wiersma,⁴ and F.W. Wise²

¹*Department of Chemistry and Chemical Biology, Cornell University, Ithaca, NY 14853, USA*

²*Department of Applied and Engineering Physics, Cornell University, Ithaca, NY 14853, USA*

³*Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, UK*

⁴*European Laboratory for Non-linear Spectroscopy and INFM,*

Via Nello Carrara 1, 50019 Sesto-Fiorentino, Italy

(Dated: February 2, 2008)

We report the observation of sum-frequency signals that depend linearly on an applied electrostatic field and that change sign with the handedness of an optically active solution. This recently predicted chiral electro-optic effect exists in the electric-dipole approximation. The static electric field gives rise to an electric-field-induced sum-frequency signal (an achiral third-order process) that interferes with the chirality-specific sum-frequency at second-order. The cross-terms linear in the electrostatic field constitute the effect and may be used to determine the absolute sign of second- and third-order nonlinear optical susceptibilities in isotropic media.

Most biological molecules are chiral [1], i.e. have a non-superimposable mirror image. In many respects the two mirror image forms (enantiomers) of a chiral molecule have identical physical properties. Optical activity is often the only practical means to distinguish between enantiomers in solution. Conventional optical activity phenomena, such as optical rotation and circular dichroism, are based on the interference of induced oscillating electric- and magnetic (and electric-quadrupole) moments, and arise from a differential response to left and right circularly polarized light [2].

Remarkably, a purely electric-dipolar nonlinear-optical process can also be a probe of chirality [3]. Two optical fields of different frequency can coherently mix in an isotropic medium to generate light at their sum (or difference) frequency if the medium is optically active, i.e. a non-racemic solution of chiral molecules [3]. The signal itself is then a probe of molecular chirality: It is the intrinsic broken symmetry of the chiral molecules that causes a non-racemic liquid to be noncentrosymmetric and hence allows for an electric-dipolar second-order nonlinear optical process in an isotropic medium.

The sum-frequency intensity is in general, however, not sensitive to the sign of the underlying property tensor, and thus does not readily distinguish between enantiomers. In this Letter we report the observation of a recently predicted chiral electro-optic effect [4] that arises when a static electric field is applied to coherent sum- or difference-frequency generation in an optically active liquid. The static field does not change the phase matching conditions, but it gives rise to an electric-field-induced contribution to the signal. The beat between chirality-sensitive sum-frequency generation (a second-order process) and achiral electric-field-induced sum-frequency generation (a third order process) yields a contribution to the intensity that is linear in the static electric field and that changes sign with the enantiomer [4, 5]. The effect can give the absolute sign of the

isotropic part of the sum-frequency hyperpolarizability (if the sign of the achiral third-order property is known) and hence makes it possible to determine the handedness of chiral molecules in solution via an electric-dipolar optical process.

Sum-frequency generation (SFG) from chiral liquids has recently been re-examined [6] and has been observed experimentally [7, 8, 9]. The molecular response to two optical fields $E_\beta(\omega_1)$ and $E_\gamma(\omega_2)$ (and a static field $E_\delta(0)$) at the sum-frequency $\omega_3 = \omega_1 + \omega_2$ can be written in terms of an induced oscillating dipole moment

$$\begin{aligned} \mu_\alpha(\omega_3) = & \frac{1}{2} \beta_{\alpha\beta\gamma}(\omega_3 = \omega_1 + \omega_2) E_\beta(\omega_1) E_\gamma(\omega_2) \quad (1) \\ & + \frac{1}{2} \gamma_{\alpha\beta\gamma\delta}(\omega_3 = \omega_1 + \omega_2 + 0) E_\beta(\omega_1) E_\gamma(\omega_2) E_\delta(0), \end{aligned}$$

where $\beta_{\alpha\beta\gamma}$ and $\gamma_{\alpha\beta\gamma\delta}$ are the first and second hyperpolarizability respectively. The macroscopic response is given by an appropriate spatial average. Here we consider an isotropic medium in the presence of a static electric field $E_\delta(0)$. A Boltzmann average yields the induced macroscopic polarization at the sum-frequency [4, 5]:

$$\begin{aligned} P_\alpha(\omega_3) = & \epsilon_0 E_\beta(\omega_1) E_\gamma(\omega_2) \left[\overbrace{\epsilon_{\alpha\beta\gamma} \chi^{(2)}}^{\text{chiral}} \right. \quad (2) \\ & \left. + 3 \underbrace{\left(\chi_1^{(3)} \delta_{\alpha\beta} \delta_{\gamma\delta} + \chi_2^{(3)} \delta_{\alpha\gamma} \delta_{\beta\delta} + \chi_3^{(3)} \delta_{\alpha\delta} \delta_{\beta\gamma} \right) E_\delta(0)}_{\text{achiral}} \right], \end{aligned}$$

where $\epsilon_{\alpha\beta\gamma}$ is the Levi-Civita tensor and $\delta_{\alpha\beta}$ the Kronecker delta. $\chi^{(2)}$ is, as we will show, the completely antisymmetric chirally sensitive isotropic component of the second-order susceptibility, and $\chi_i^{(3)}$ with $i = 1, 2$ or 3 are the achiral isotropic components of the third-order susceptibility. The intensity at the sum-frequency is proportional to $|P_\alpha(\omega_3)|^2$ and so contains – besides a contribution independent of the static field, and one that is

quadratic in the static field – cross terms that are linear in the static field and linear in $\chi^{(2)}$:

$$\text{SFG}(E) \propto \text{Re} \left[\chi^{(2)} \left(\chi_i^{(3)} \right)^* \right] E(0) I(\omega_1) I(\omega_2), \quad (3)$$

where I are the respective incident intensities, Re is the real part and the star indicates a complex conjugate. We have denoted the contribution to the intensity at the sum-frequency that is linear in the static field by $\text{SFG}(E)$.

$\chi^{(2)}$ is proportional to the isotropic component of the first hyperpolarizability and is given by

$$\chi^{(2)} = \frac{N}{12 \epsilon_0} (\beta_{xyz} - \beta_{xzy} + \beta_{yzx} - \beta_{yxz} + \beta_{zxy} - \beta_{zyx}),$$

where N is the excess number density. It is seen that $\chi^{(2)}$ vanishes for any molecule that possesses a mirror plane, a center of inversion, or a rotation-reflection axis; thus $\chi^{(2)}$ is only non-zero for chiral systems and changes sign as the handedness of the optically active liquid changes. The contribution to the intensity linear in $E(0)$ may therefore reveal the sign of the pseudoscalar $\chi^{(2)}$. It can be used to determine the absolute configuration of the chiral molecules in the optically active liquid.

$\chi_1^{(3)}$, $\chi_2^{(3)}$ and $\chi_3^{(3)}$ are given in terms of scalar combinations of the second hyperpolarizability tensor components by [4]:

$$\begin{aligned} \chi_1^{(3)} &= \frac{N}{180 \epsilon_0} [4\gamma_{\alpha\alpha\beta\beta} - \gamma_{\alpha\beta\alpha\beta} - \gamma_{\alpha\beta\beta\alpha}] \\ \chi_2^{(3)} &= \frac{N}{180 \epsilon_0} [-\gamma_{\alpha\alpha\beta\beta} + 4\gamma_{\alpha\beta\alpha\beta} - \gamma_{\alpha\beta\beta\alpha}] \\ \chi_3^{(3)} &= \frac{N}{180 \epsilon_0} [-\gamma_{\alpha\alpha\beta\beta} - \gamma_{\alpha\beta\alpha\beta} + 4\gamma_{\alpha\beta\beta\alpha}]. \end{aligned} \quad (4)$$

Should the duration of the static applied field be longer than the rotational time of the molecules in solution, then there are additional temperature-dependent terms of the form $\beta_{\alpha\alpha\beta}(\omega_3 = \omega_1 + \omega_2) \mu_\beta^{(0)} / (kT)$ for molecules with a permanent dipole moment $\mu_\beta^{(0)}$ [4]. At ambient conditions, the temperature dependent contributions may dominate for dipolar molecules and be of either sign. Both $\gamma_{\alpha\alpha\beta\beta}$ and $\beta_{\alpha\alpha\beta} \mu_\beta^{(0)}$ are unchanged under mirror symmetry operation and so it follows that the $\chi_i^{(3)}$ exist for achiral molecules (e.g. an achiral solvent) and are necessarily the same for both enantiomers of a chiral molecule.

We observe the chiral electro-optic effect in an experimental arrangement schematically depicted in Figure 1a. From the antisymmetry of the Levi-Civita tensor in Eqn. (2) it follows that the optical fields need to have components that are mutually orthogonal such that they span the x , y , and z axes of a Cartesian frame. Hence, sum-frequency generation from isotropic media requires that one optical field be S-polarized and the remaining two be P-polarized. Further, the static electric field has to be

S-polarized for it to give rise to an electric-field-induced contribution to the sum-frequency signal.

We use the 775-nm fundamental wavelength of a Ti:sapphire regenerative amplifier (Clark-MXR CPA-2010) along with its second harmonic at 338 nm to generate sum-frequency signals at 258 nm. The pulse duration is ~ 150 fs and the repetition rate is 1 kHz. A custom-built high voltage power supply provides 3.5 to 10 kV peak to peak modulated at 500 Hz. The modulation of the applied electric field permits phase sensitive detection.

Unlike in collinear electric-field-induced second-harmonic generation (EFISH) [10], the non-collinear SFG beam geometry [6] allows the use of fully immersed electrodes. These are suspended in a standard quartz optical cuvette and spaced ~ 2 mm apart.

We observe the chiral electro-optic effect in optically active solutions prepared from R-(+)- or S-(−)-1, 1'-bi-2-naphthol (R-BN and S-BN) (see Figure 1b) which are dissolved in the achiral solvent tetrahydrofuran (THF). In Figure 2 we show that the signals are of opposite sign for the R- and S-enantiomers of BN and that they vary linearly with the strength of the applied low-frequency field. The small difference in absolute strength of the respective signals is attributed to small differences in the electrode-spacings and/or the concentrations. The $\text{SFG}(E)$ signals change sign as the direction of the static field is reversed.

For fixed beam polarizations, direction and strength of the static field, we also observe a linear dependence of the chiral electro-optic effect on the enantiomeric concentration difference of the optically active solution, as is seen in Figure 3. Starting with a 0.56 M solution of R-(+)-1, 1'-bi-2-naphthol in THF, the handedness of the 0.56 M solution is gradually changed by the addition of a solution of S-(−)-1, 1'-bi-2-naphthol in THF. The $\text{SFG}(E)$ signal correspondingly changes sign. It is zero for the racemic mixture.

To describe the polarization dependence of the sum-frequency signals, we now consider the axis system and beam geometry depicted in Figure 1a. For simplicity we ignore the small rotation of the plane of polarization in the chiral liquid due to optical activity ($< 0.5^\circ$), and we note that a static electric field can not contribute linearly to optical rotation in a fluid.

From Eqn. (2) and the requirement for a transverse wave at the sum-frequency, it follows that a PPS combination of polarizations (listed throughout this Letter in the order of the optical fields at: $\omega_3, \omega_1, \omega_2$) gives rise to an $\text{SFG}(E)$ signal of the form

$$\text{SFG}(E)_{\text{PPS}} = (+) \eta \text{Re} \left[\chi^{(2)} \left(\chi_1^{(3)} \right)^* \right] \sin(2\theta_1) E_x(0),$$

where θ_1 is the angle the incident beam at ω_1 makes with respect to the sum-frequency beam, and where we have subsumed the incident intensities and any numerical factors common to all $\text{SFG}(E)$ intensities in the positive

multiplier η . Similarly, PSP polarizations give rise to

$$\text{SFG(E)}_{\text{PSP}} = (+) \eta \text{Re} \left[\chi^{(2)} \left(\chi_3^{(3)} \right)^* \right] \sin(2\theta_2) E_x(0),$$

and SPP polarizations probe

$$\text{SFG(E)}_{\text{SPP}} = (-) \eta \text{Re} \left[\chi^{(2)} \left(\chi_3^{(3)} \right)^* \right] \times \sin(2(\theta_1 + \theta_2)) E_x(0).$$

Since $\chi_i^{(3)}$ are the same for the R- and S-enantiomers while the $\chi^{(2)}$ change sign, solutions of R- and S-BN with equal concentration will have equal but opposite SFG(E) intensities in the same field $E_x(0)$, and this can be seen in Figure 4. Further, the change of sign in going from PPS to SPP polarizations and similarly from SPP to PSP suggests that $\chi_1^{(3)}$, $\chi_2^{(3)}$ and $\chi_3^{(3)}$ have the same sign for BN ($2(\theta_1 + \theta_2) < \pi$). The three polarization states PPS, PSP and SPP permit the observation of the three isotropic third-order susceptibilities, $\chi_i^{(3)}$. For a static field of ~ 2.5 kV/cm the SFG(E) signals are, depending on beam polarizations, ~ 5 to 20% of the SFG signals in the absence of a dc electric field shown in Figure 5. Nonlinear optical susceptibilities of different order may thus be measured under identical optical field conditions without the need for an external reference, as the $\chi^{(2)}$ will act as an internal standard. The relative strength of $\chi_1^{(3)}$, $\chi_2^{(3)}$ and $\chi_3^{(3)}$ may in turn yield information about specific hyperpolarizability tensor components entering the third-order susceptibility. Such an analysis is, however, chromophore specific and will not be discussed further here. Finally, Figure 5 confirms that SFG signals independent of E do not distinguish between enantiomers.

For a dc electric field along $+x$ (see Fig 1a) we measure a positive $\text{SFG(E)}_{\text{PSP}}$ intensity for S-BN. In the case of the resonant signals from R- and S-BN, where the complex nature of the response tensors needs to be considered, and in the absence of any measurable $\chi_i^{(3)}$ contribution from the achiral solvent, the determination of the absolute configuration of the chiral molecules will require quantum chemical computations for both $\chi^{(2)}$ and $\chi_i^{(3)}$. However, the contribution from an achiral solvent should make it possible to determine the absolute sign of both the second- and third-order susceptibilities for both optical isomers of a chiral molecule [4].

In summary, we have observed the linear effect of a static electric field on the sum-frequency generation intensity from optically active liquids. The reported chiral electro-optic effect arises in the electric-dipole approximation and changes sign with the enantiomer and upon reversing the direction of the static electric field. The effect may be used to determine the absolute sign of the second-order and third-order nonlinear optical susceptibilities.

Should the achiral solvent give rise to an appreciable electric-field-induced sum-frequency signal, then the ef-

fect could be used to amplify weak sum-frequency signals from chiral solutions.

The authors are grateful to Professor A.C. Albrecht for many helpful discussions, and thank Yi-Fan Chen for assistance with the experiment. Clark-MXR is acknowledged for the use of the regenerative amplifier. This work was supported by the National Science Foundation (CHE-0095056) and the National Institutes of Health (EB002019).

* Corresponding author: pf43@cornell.edu

- [1] L. Kelvin, *Baltimore Lectures* (C.J. Clay & Sons, London, 1904.)
- [2] L. D. Barron, *Molecular Light Scattering and Optical Activity* (CUP, Cambridge, 1982).
- [3] J. A. Giordmaine, Phys. Rev. **138**, A1599 (1965).
- [4] A. D. Buckingham and P. Fischer, Chem. Phys. Letters **297**, 239 (1998).
- [5] A. D. Buckingham and P. Fischer, in *Physical Chemistry of Chirality*, edited by J. M. Hicks (Oxford University Press, Oxford, 2002), p. 119.
- [6] P. Fischer, D. S. Wiersma, R. Righini, B. Champagne, and A. D. Buckingham, Phys. Rev. Letters **85**, 4253 (2000).
- [7] M. A. Belkin, T. A. Kulakov, K.-H. Ernst, L. Yan, and Y. R. Shen, Phys. Rev. Letters **85**, 4474 (2000).
- [8] M. A. Belkin, S. H. Han, X. Wei, and Y. R. Shen, Phys. Rev. Letters **87**, 113001 (2001).
- [9] P. Fischer, K. Beckwitt, F. W. Wise, and A. C. Albrecht, Chem. Phys. Letters **352**, 463 (2002).
- [10] B. F. Levine and C. G. Bethea, J. Chem. Phys. **63**, 2666 (1975).




FIG. 1: (color online) a) Beam geometry and choice of axes. The optical beams lie in the y, z plane, and the static field is S-polarized along x . The incident beam at ω_1 makes an angle of θ_1 with the sum-frequency beam at ω_3 ; and similarly for ω_2 . (The cuvette with the optically active liquid is not shown.) b) Structure of R-(+)-1,1'-bi-2-naphthol ($\text{C}_{20}\text{H}_{14}\text{O}_2$).




FIG. 2: (color online) SFG(E) signals from ~ 0.5 M solutions of S-($-$)- and R-($+$)-1,1'-bi-2-naphthol respectively measured at different strengths (estimated) of the applied electric field. The solid lines are linear functions fitted to the data.




FIG. 3: SFG(E) signals measured as a function of the fractional concentration difference of R-($+$)- and S-($-$)-1,1'-bi-2-naphthol in THF. The solid line is a linear function fitted to the data.




FIG. 4: (color online) Sum-frequency generation signals from optically active ~ 0.5 M solutions of S(-)- and R(+)-BN in THF observed for different beam polarizations. The polarization state of the ω_1 beam is shown on the lower axis and that of the ω_2 beam on the upper axis. First, $E(\omega_1)$ is kept P-polarized while the polarization of $E(\omega_2)$ is changed from S to P in 9° increments. Subsequently, $E(\omega_2)$ remains P-polarized while the polarization of the ω_1 beam is changed from P to S. When both the incident optical fields at ω_1 and ω_2 are P-polarized, then the ω_3 beam is S-polarized. Mixed input polarizations (S and P) give rise to a P-polarized signal. The static electric field E_x is ~ 2 kV/cm. The expected trend of the signals is shown by a model calculation depicted in the inset.




FIG. 5: (color online) The SFG intensity in the absence of a static electric field is shown as a function of the input beam polarizations (see caption Fig. 4 for details). The solid line is a fit from theory. The contribution quadratic in the static field is below the sensitivity of the experiment.